

hCCR8(2)/hCTLA-4

品系全名	C57BL/6Smoc- <i>Ccr8</i> ^{em3(hCCR8)} <i>Ctla4</i> ^{em1(hCTLA4)Smoc}
目录号	NM-HU-220074
品系状态	活体

基因信息

基因名 CCR8	基因曾用名	Cmkbr8
	NCBI ID	12776
	MGI ID	1201402
	Ensembl ID	ENSMUSG00000042262
	人类同源基因	CCR8
基因名 CTLA4	基因曾用名	Cd152; Ly-56; Ctla-4
	NCBI ID	12477
	MGI ID	88556
	Ensembl ID	ENSMUSG00000026011
	人类同源基因	CTLA4

品系描述

通过hCTLA4 (NM-HU-00014) 与hCCR8(2) (NM-HU-2000054) 交配获得hCCR8(2)/hCTLA4小鼠。

*使用本品系发表的文献需注明: hCCR8(2)/hCTLA-4 mice (Cat. NO. NM-HU-220074) were purchased from Shanghai Model Organisms Center, Inc..

验证数据




Fig1. Statistics of hCCR8 and hCTLA-4 expression in hCCR8(2)/hCTLA-4 dKI mice.




Fig2. Detection of hCCR8 and CTLA-4 expression on T cells in Tumor in hCCR8(2)/hCTLA-4 dKI mice.




Fig3. Detection of hCCR8 and hCTLA-4 expression on T cells in Spleen in hCCR8(2)/hCTLA-4 dKI mice.

